

An algorithm for solving the variational inequality problem over the fixed point set of a quasi-nonexpansive operator in Euclidean space

Andrzej Cegielski¹, Aviv Gibali², Simeon Reich² and Rafał Zalas¹

¹Faculty of Mathematics, Computer Science and Econometrics,
University of Zielona Góra, Zielona Góra, Poland

²Department of Mathematics,
The Technion - Israel Institute of Technology
Technion City, 32000 Haifa, Israel

July 25, 2012, Revised December 17, 2012.

Abstract

This paper is concerned with the variational inequality problem $VIP(\mathcal{F}, \text{Fix}(T))$: find $\bar{u} \in \text{Fix}(T)$ such that $\langle \mathcal{F}(\bar{u}), z - \bar{u} \rangle \geq 0$ for all $z \in \text{Fix}(T)$, where $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is quasi-nonexpansive, $\text{Fix}(T)$ is its nonempty fixed point set, and $\mathcal{F} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is monotone. We propose, in particular, an algorithm which entails, at each step, projecting onto a suitably chosen half-space, and prove that the sequences it generates converge to the unique solution of the VIP. We also present an application of our result to a hierarchical optimization problem.

Key words: Fixed point, quasi-nonexpansive operator, variational inequality problem.

AMS Mathematical Subject Classification: 47H05, 47H09, 47H10, 47J20, 47J40, 65K15, 90C23.

1 Introduction

The classical variational inequality problem (VIP) is to find a point $x^* \in S$ such that

$$\langle \mathcal{F}(x^*), x - x^* \rangle \geq 0 \text{ for all } x \in S, \quad (1.1)$$

where $S \subseteq \mathbb{R}^n$ is nonempty, closed and convex, $\mathcal{F} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a given operator, and $\langle \cdot, \cdot \rangle$ denotes the inner product in $\mathbb{R}^n$. This problem, denoted by $\text{VIP}(\mathcal{F}, S)$, is a fundamental problem in Optimization Theory because many optimization problems can be translated into VIPs. The VIP was intensively studied in the last decades; see, *e.g.*, the two-volume book by Facchinei and Pang [20], and the review papers by Noor [30] and by Xiu and Zhang [32]. Some algorithms for solving (1.1) fit into the framework of the following general iterative scheme:

$$x^{k+1} = P_S(x^k - \tau_k \mathcal{F}(x^k)), \quad (1.2)$$

where $\tau_k \geq 0$ and P_S is the metric projection operator onto S (see Auslender [2] and consult [20, Volume 2, Subsection 12.1] for more details).

Such methods are particularly useful when the set S is simple enough to project onto. However, in general, one has to solve a minimization problem (evaluation of the metric projection onto S) at each iterative step in order to get the next iterate. In this case the efficiency of method (1.2) may be seriously affected. Fukushima [22] developed a method that overcomes this obstacle by replacing the metric projection onto the set S by a subgradient projection which is easier to calculate. Censor and Gibali present in [14] the δ -algorithmic scheme which generalizes the Auslender and Fukushima algorithms in the sense that there is some “freedom” in choosing the hyperplane onto which one projects.

In this paper we present a natural extension of this δ -algorithmic scheme (Algorithm 3.7 below). Given an operator $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$, we denote by $\text{Fix}(T) := \{x \in \mathbb{R}^n \mid T(x) = x\}$ the fixed point set of T . It is well known that $\text{Fix}(T)$ is closed and convex if T is quasi-nonexpansive (see, *e.g.*, [5, Proposition 2.6 (ii)]). Observe that the feasible set S of the VIP in (1.1) can always be represented as the fixed point set of some operator, say, $S = \text{Fix}(P_S)$. Following this idea, Yamada and Ogura [34] considered the variational inequality problem $\text{VIP}(\mathcal{F}, \text{Fix}(T))$, which calls for finding a point $x^* \in \text{Fix}(T)$ such that

$$\langle \mathcal{F}(x^*), x - x^* \rangle \geq 0 \text{ for all } x \in \text{Fix}(T). \quad (1.3)$$

In the case where T is quasi-nonexpansive and so-called quasi-shrinking, an algorithm for solving (1.3) in a real Hilbert space was proposed in [34] under the conditions of Lipschitz continuity and strong monotonicity of $\mathcal{F}$. The iterative step of the method is as follows:

$$x^{k+1} = T(x^k) - \lambda_{k+1}\mathcal{F}(T(x^k)), \quad (1.4)$$

where $\{\lambda_k\}_{k=0}^{\infty}$ is a nonnegative sequence which satisfies certain conditions. As a matter of fact, Yamada and Ogura [34, Theorem 5] showed that (1.4) could be applied to more general cases with weaker monotonicity assumptions, such as paramonotonicity (see e.g., [35]).

In this paper we present a method for solving the $\text{VIP}(\mathcal{F}, \text{Fix}(T))$ when the operator T is merely quasi-nonexpansive. This method generalizes the earlier results of Auslender and Fukushima, as well as the δ -algorithmic scheme. In addition, we present the relationship between our algorithm and the Yamada–Ogura method. Note that several authors have considered the $\text{VIP}(\mathcal{F}, \text{Fix}(T))$ for a quasi-nonexpansive operator T and proposed methods similar to (1.4), where T is replaced by a sequence of quasi-nonexpansive operators T_k with the property $\text{Fix}(T) \subseteq \bigcap_{k \geq 0} \text{Fix}(T_k)$ (see [3, 13, 26, 34]).

Our paper is organized as follows. Section 2 is devoted to some definitions and preliminary results. Our algorithm is described in Section 3 and analyzed in Section 4. Finally, in Section 5 we present an application of our results to a hierarchical optimization problem.

2 Preliminaries

In this section we recall some definitions and properties of several classes of operators.

Definition 2.1 Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be an operator with a fixed point. The operator T is called:

- (i) α -strongly quasi-nonexpansive (α -SQNE), where $\alpha \geq 0$, if for all $(x, w) \in \mathbb{R}^n \times \text{Fix}(T)$, we have

$$\|T(x) - w\|^2 \leq \|x - w\|^2 - \alpha \|x - T(x)\|^2; \quad (2.1)$$

If $\alpha > 0$, then we say that T is strongly quasi-nonexpansive (SQNE);

- (ii) **Firmly quasi-nonexpansive** (FQNE) if it is 1-SQNE;
- (iii) **Quasi-nonexpansive** if it is 0-SQNE, i.e.,

$$\|T(x) - w\| \leq \|x - w\| \text{ for all } (x, w) \in \mathbb{R}^n \times \text{Fix}(T); \quad (2.2)$$

- (iv) **Nonexpansive** if

$$\|T(x) - T(y)\| \leq \|x - y\| \text{ for all } x, y \in \mathbb{R}^n. \quad (2.3)$$

The class of quasi-nonexpansive operators was denoted by Crombez [18, p. 161] by $\mathcal{F}^0$. An important subset of $\mathcal{F}^0$, namely the $\mathfrak{T}$ -class operators, was introduced and investigated by Bauschke and Combettes [5], and by Combettes [17]. The operators in this class were named *directed operators* in Zaknoon [36] and further used under this name in [16]. Cegielski [10] studied these operators under the name *separating operators*. Since both *directed* and *separating* are key words of other, widely-used, mathematical entities, Cegielski and Censor have recently introduced the term *cutter operators* [12], or cutters in short. This class coincides with the class $\mathcal{F}^1$ (see [18]), with the class of 1-SQNE operators (see [11, Theorem 2.1.39]) and with the class $\text{DC}_{\mathbf{p}}$ for $\mathbf{p} = -1$ [29]. The term *firmly quasi-nonexpansive* (FQNE) for $\mathfrak{T}$ -class operators was used by Yamada and Ogura [33, 34, Section B] and by Mărușter [28]. This class of operators is fundamental because it contains several types of operators commonly found in various areas of applied mathematics, such as the metric projections, subgradient projections and the resolvents of maximal monotone operators (see [5]). The formal definition of the $\mathfrak{T}$ -class in Euclidean space is as follows.

Definition 2.2 An operator $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is called a **cutter** ($T \in \mathfrak{T}$) if $\text{Fix}(T) \neq \emptyset$ and

$$\langle T(x) - x, T(x) - w \rangle \leq 0 \text{ for all } (x, w) \in \mathbb{R}^n \times \text{Fix}(T). \quad (2.4)$$

For $x, y \in \mathbb{R}^n$, we denote

$$H(x, y) := \{u \in \mathbb{R}^n \mid \langle u - y, x - y \rangle \leq 0\}. \quad (2.5)$$

If $x \neq y$, then $H(x, y)$ is a half-space. It is easy to see that T is a cutter if and only if

$$\text{Fix}(T) \subseteq H(x, T(x)) \text{ for all } x \in \mathbb{R}^n. \quad (2.6)$$

This property is illustrated in Figure 1.




Figure 1: Property of cutters

Definition 2.3 Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a cutter and $\alpha \in [0, 2]$. The operator $T_\alpha := I + \alpha(T - I)$, where I denotes the identity operator, is called an α -relaxed cutter.

Bauschke and Combettes [5] established the following two properties of cutters.

- (i) The set $\text{Fix}(T)$ of all fixed points of a cutter operator T is closed and convex because

$$\text{Fix}(T) = \bigcap_{x \in \mathbb{R}^n} H(x, T(x)). \quad (2.7)$$

- (ii) If $T \in \mathfrak{T}$ and $\alpha \in [0, 1]$, then $T_\alpha \in \mathfrak{T}$.

One can easily verify the following characterization of α -relaxed cutter operators U ($U = T_\alpha$):

$$\alpha \langle U(x) - x, w - x \rangle \geq \|U(x) - x\|^2 \text{ for all } (x, w) \in \mathbb{R}^n \times \text{Fix}(T). \quad (2.8)$$

Theorem 2.4 Let $\alpha \in (0, 2]$. An operator $U : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a cutter if and only if U_α is $((2 - \alpha) / \alpha)$ -strongly quasi-nonexpansive

Proof. See, e.g., [17, Proposition 2.3 (ii)] and [11, Theorem 2.1.39]. ■

Notation 2.5 Let C be a nonempty, closed and convex subset of $\mathbb{R}^n$, and let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be quasi-nonexpansive.

(i) The **distance function** of a point $x \in \mathbb{R}^n$ to C is defined by

$$\text{dist}(x, C) := \inf\{\|x - z\| \mid z \in C\}. \quad (2.9)$$

(ii) For $r \geq 0$ we define the subsets

$$C^r := \{u \in \mathbb{R}^n \mid \text{dist}(u, C) \geq r\} \quad (2.10)$$

and

$$C_r := \{u \in \mathbb{R}^n \mid \text{dist}(u, C) \leq r\}. \quad (2.11)$$

(iii) Define the function $D: [0, \infty) \rightarrow [0, \infty]$ in the following way:

$$D(r) := \begin{cases} \inf_{u \in (\text{Fix}(T))^r \cap C} (\text{dist}(u, \text{Fix}(T)) - \text{dist}(T(u), \text{Fix}(T))) \\ \text{if } (\text{Fix}(T))^r \cap C \neq \emptyset, \\ +\infty, \text{ otherwise.} \end{cases} \quad (2.12)$$

(iv) Denote the metric projection onto $\text{Fix}(T)$ by R , i.e., $R := P_{\text{Fix}(T)}$.

It is well known that, for a convex subset C , the (continuous) distance function $\text{dist}(\cdot, C)$ is convex. Consequently, C_r is closed and convex as a sublevel set of a convex function. The continuity of $\text{dist}(\cdot, C)$ implies that the subset C^r is also closed.

In what follows we assume that $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is quasi-nonexpansive and $C \subseteq \mathbb{R}^n$ is a closed and convex set such that $\text{Fix}(T) \cap C \neq \emptyset$.

Parts (i) and (ii) of the following theorem can be found in [34, Lemma 2].

Proposition 2.6 *The function $D: [0, \infty) \rightarrow [0, \infty]$ defined by (2.12) has the following properties:*

- (i) $D(0) = 0$ and $D(r) \geq 0$ for all $r \geq 0$;
- (ii) if $r_1 \geq r_2 \geq 0$, then $D(r_1) \geq D(r_2)$;
- (iii) $D(\text{dist}(x, \text{Fix}(T))) \leq \|x - T(x)\|$ for all $x \in C$.

Proof. (i) Let $r \geq 0$. We prove that $D(r) \geq 0$. The inequality is clear if $(\text{Fix}(T))^r \cap C = \emptyset$. Now suppose that $(\text{Fix}(T))^r \cap C \neq \emptyset$. Then the definition of the metric projection and the quasi-nonexpansivity of T yield, for any $x \in (\text{Fix}(T))^r \cap C$,

$$\|x - R(x)\| - \|T(x) - R(T(x))\| \geq \|x - R(x)\| - \|T(x) - R(x)\| \geq 0. \quad (2.13)$$

Consequently, $D(r) \geq 0$. Let $x \in \text{Fix}(T) \cap C$. By the quasi-nonexpansivity of T , we have $T(x) = x$ and

$$D(0) \leq \|x - R(x)\| - \|T(x) - R(T(x))\| = 0, \quad (2.14)$$

which together with the first part proves that $D(0) = 0$.

(ii) Let $r_1 \geq r_2 \geq 0$. Then, of course, $(\text{Fix}(T))^{r_2} \subseteq (\text{Fix}(T))^{r_1}$ and so the property is clear.

(iii) Let $x \in C$ and $r = \text{dist}(x, \text{Fix}(T))$. If $r = 0$, then $T(x) = x$ and, by (i), the assertion is obvious. Let $r > 0$. Then, of course, $x \in (\text{Fix}(T))^r \cap C$ and, by the definition of the metric projection and the triangle inequality, we have

$$\begin{aligned} D(r) &\leq \|x - R(x)\| - \|T(x) - R(T(x))\| \\ &\leq \|x - R(T(x))\| - \|T(x) - R(T(x))\| \leq \|x - T(x)\|, \end{aligned} \quad (2.15)$$

and the proof is complete. ■

Now we give two equivalent definitions of a quasi-shrinking operator.

Definition 2.7 (cf. [34]) Let $C \subseteq \mathbb{R}^n$ be closed and convex, and let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a quasi-nonexpansive operator. The operator T is called **quasi-shrinking** on C if $D(r) = 0 \Leftrightarrow r = 0$.

Definition 2.8 Let $C \subseteq \mathbb{R}^n$ be closed and convex, and let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a quasi-nonexpansive operator. We say that T is **quasi-shrinking** on C if for any sequence $\{u^k\}_{k=0}^\infty \subseteq C$, the following implication holds:

$$\lim_{k \rightarrow \infty} (\|u^k - R(u^k)\| - \|T(u^k) - R(T(u^k))\|) = 0 \Rightarrow \lim_{k \rightarrow \infty} \|u^k - R(u^k)\| = 0. \quad (2.16)$$

Proposition 2.9 Let $C \subseteq \mathbb{R}^n$ be closed and convex, and let a quasi-nonexpansive operator $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be such that $\text{Fix}(T) \cap C \neq \emptyset$. Then Definitions 2.7 and 2.8 are equivalent.

Proof. Let T be quasi-shrinking in the sense of Definition 2.7 and choose $\{u^k\}_{k=0}^\infty \subseteq C$. Suppose that $\lim_{k \rightarrow \infty} \|u^k - R(u^k)\| \neq 0$. Then there exist a constant $\varepsilon > 0$ and a subsequence $\{u^{k_j}\}_{j=0}^\infty \subseteq \{u^k\}_{k=0}^\infty$ such that $\|u^{k_j} - R(u^{k_j})\| > \varepsilon$. Therefore $u^{k_j} \in (\text{Fix}(T))^\varepsilon \cap C$ and we have

$$\begin{aligned} & \inf_{j \geq 0} (\|u^{k_j} - R(u^{k_j})\| - \|T(u^{k_j}) - R(T(u^{k_j}))\|) \\ & \geq \inf_{u \in (\text{Fix}(T))^\varepsilon \cap C} (\|u - R(u)\| - \|T(u) - R(T(u))\|) \\ & = D(\varepsilon) > 0. \end{aligned} \quad (2.17)$$

Consequently, $\lim_{k \rightarrow \infty} (\|u^k - R(u^k)\| - \|T(u^k) - R(T(u^k))\|) \neq 0$ if it exists.

In the other direction, let T be quasi-shrinking in the sense of Definition 2.8. Suppose that $D(r) = 0$ for some $r \geq 0$. Then there is a sequence $\{u^k\}_{k=0}^\infty \subseteq (\text{Fix}(T))^r \cap C$ such that

$$\lim_{k \rightarrow \infty} (\|u^k - R(u^k)\| - \|T(u^k) - R(T(u^k))\|) = 0. \quad (2.18)$$

By Definition 2.8, we have

$$r \leq \lim_{k \rightarrow \infty} \text{dist}(u^k, \text{Fix}(T)) = \lim_{k \rightarrow \infty} \|u^k - R(u^k)\| = 0, \quad (2.19)$$

i.e., $r = 0$ and the proof is complete. ■

Definition 2.10 Let $C \subseteq \mathbb{R}^n$ be closed and convex. An operator $T : C \rightarrow \mathbb{R}^n$ is called **closed** at $y \in \mathbb{R}^n$ if for any sequence $\{x^k\}_{k=0}^\infty \subseteq C$, we have

$$\left(\lim_{k \rightarrow \infty} x^k = x \in C \text{ and } \lim_{k \rightarrow \infty} T(x^k) = y \right) \Rightarrow T(x) = y. \quad (2.20)$$

We say that the *closedness principle* holds for an operator $T : C \rightarrow \mathbb{R}^n$ if $I - T$ is closed at 0 (see [8]), i.e.,

$$\left(\lim_{k \rightarrow \infty} x^k = x \in C \text{ and } \lim_{k \rightarrow \infty} \|T(x^k) - x^k\| = 0 \right) \Rightarrow x \in \text{Fix}(T). \quad (2.21)$$

It is clear that in $\mathbb{R}^n$ a continuous operator, in particular a nonexpansive one, satisfies the closedness principle. Later in this section we give other examples of operators satisfying the closedness principle (see Examples 2.14, 2.19 and 2.22).

Proposition 2.11 *Let $C \subseteq \mathbb{R}^n$ be closed, bounded and convex and let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be an operator with $\text{Fix}(T) \cap C \neq \emptyset$. If T is SQNE (equivalently, an α -relaxed cutter for some $\alpha \in (0, 2)$) and $I - T$ is closed at 0, then T is quasi-shrinking on C .*

Proof. Let $r \geq 0$ and $D(r) = 0$. Then there is a sequence $\{u^k\}_{k=0}^\infty \subseteq (\text{Fix}(T))^r \cap C$ such that

$$\lim_{k \rightarrow \infty} (\|u^k - R(u^k)\| - \|T(u^k) - R(T(u^k))\|) = 0. \quad (2.22)$$

By the quasi-nonexpansivity of T , the definition of the metric projection and by (2.22), we have

$$\begin{aligned} 0 &\leq \|u^k - R(u^k)\| - \|T(u^k) - R(u^k)\| \\ &\leq \|u^k - R(u^k)\| - \|T(u^k) - R(T(u^k))\| \rightarrow 0. \end{aligned} \quad (2.23)$$

Consequently,

$$0 \leq \|u^k - R(u^k)\| - \|T(u^k) - R(u^k)\| \rightarrow 0. \quad (2.24)$$

Since T is SQNE, there is $\alpha > 0$ such that

$$\|T(u^k) - R(u^k)\|^2 \leq \|u^k - R(u^k)\|^2 - \alpha \|T(u^k) - u^k\|^2. \quad (2.25)$$

Let $z \in \text{Fix}(T)$. By the boundedness of $\{u^k\}_{k=0}^\infty$, there exists $d > 0$ such that $\|u^k - z\| \leq d$ for all $k \geq 0$. Using the definition of the metric projection and the quasi-nonexpansivity of T , we obtain

$$\begin{aligned} \|u^k - R(u^k)\| + \|T(u^k) - R(u^k)\| &\leq \|u^k - z\| + \|u^k - R(u^k)\| \\ &\leq 2\|u^k - z\| \leq 2d. \end{aligned} \quad (2.26)$$

By (2.24), (2.25) and (2.26), we now have

$$\begin{aligned} &\|T(u^k) - u^k\|^2 \\ &\leq \frac{1}{\alpha} (\|u^k - R(u^k)\|^2 - \|T(u^k) - R(u^k)\|^2) \\ &= \frac{1}{\alpha} (\|u^k - R(u^k)\| - \|T(u^k) - R(u^k)\|) (\|u^k - R(u^k)\| + \|T(u^k) - R(u^k)\|) \\ &\leq \frac{2d}{\alpha} (\|u^k - R(u^k)\| - \|T(u^k) - R(u^k)\|) \rightarrow 0. \end{aligned} \quad (2.27)$$

Consequently,

$$\lim_{k \rightarrow \infty} \|T(u^k) - u^k\| = 0. \quad (2.28)$$

Since $\{u^k\}_{k=0}^\infty$ is bounded, there exists a subsequence $\{u^{k_j}\}_{j=0}^\infty$ of $\{u^k\}_{k=0}^\infty$ such that

$$\lim_{j \rightarrow \infty} u^{k_j} = u^*. \quad (2.29)$$

The closedness of $I - T$ at 0 yields that $u^* \in \text{Fix}(T)$ and

$$r \leq \inf_{u \in (\text{Fix}(T))^r \cap C} \text{dist}(u, \text{Fix}(T)) \leq \lim_{j \rightarrow \infty} \|u^{k_j} - u^*\| = 0, \quad (2.30)$$

i.e., $r = 0$, which proves that T is quasi-shrinking, as asserted. ■

Remark 2.12 The converse to Proposition 2.11 is not true. To see this, take $C = \{u \in \mathbb{R}^n \mid \langle a, u \rangle \leq \beta\}$ for some $a \neq 0$, $\beta \in \mathbb{R}$ and $T = 2P_C - I$. Then T is quasi-shrinking, but T is not SQNE.

The next lemma is taken from [34, Lemma 1].

Lemma 2.13 *Let $f : [0, \infty) \rightarrow [0, \infty]$ be an increasing function such that $f(r) = 0 \Leftrightarrow r = 0$. Let $\{b_k\}_{k=0}^\infty \subseteq [0, \infty)$ be such that*

$$\lim_{k \rightarrow \infty} b_k = 0. \quad (2.31)$$

Then any sequence $\{a_k\}_{k=0}^\infty \subseteq [0, \infty)$ satisfying

$$a_{k+1} \leq a_k - f(a_k) + b_k \quad (2.32)$$

converges to 0.

Let $C \subseteq \mathbb{R}^n$ be nonempty, closed and convex. It follows from the characterization of the metric projection that P_C is a cutter. Moreover, P_C satisfies the closedness principle as a nonexpansive operator. We now present more examples of cutter operators the complements of which are closed at 0.

Example 2.14 Let $U : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be an operator with a fixed point such that $I - U$ is closed at 0 (e.g., the metric projection onto a closed and convex set), and let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a cutter such that $\text{Fix}(U) \subseteq \text{Fix}(T)$, which satisfies for any bounded sequence $\{x^k\}_{k=0}^\infty \subseteq \mathbb{R}^n$ the following implication:

$$\lim_{k \rightarrow \infty} \|T(x^k) - x^k\| = 0 \implies \lim_{k \rightarrow \infty} \|U(x^k) - x^k\| = 0 \quad (2.33)$$

(e.g., a cutter T such that $\|T(x) - x\| \geq \alpha \|U(x) - x\|$ for some constant $\alpha > 0$ and for all $x \in \mathbb{R}^n$). Then T is closed at 0.

Indeed, let $\lim_{k \rightarrow \infty} x^k = z$ and $\lim_{k \rightarrow \infty} \|T(x^k) - x^k\| = 0$. Then $\{x^k\}_{k=0}^\infty$ is bounded and, by (2.33), we have $\lim_{k \rightarrow \infty} \|U(x^k) - x^k\| = 0$. Since $I - U$ is closed at 0, we know that $z \in \text{Fix}(U)$. Consequently, $z \in \text{Fix}(T)$, i.e., $I - T$ is closed at 0. An operator T having the properties described above is a special case of the sequence of operators considered in [9, Theorem 1] and in [12, Theorem 9.9].

Next we present two more examples of operators which are cutters and the complements of which are closed at 0. These examples are special cases of Example 2.14.

First we present several definitions.

Definition 2.15 Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function.

(i) The set

$$\partial f(y) := \{g \in \mathbb{R}^n : f(x) \geq f(y) + \langle g, x - y \rangle \text{ for all } x \in \mathbb{R}^n\} \quad (2.34)$$

is called the **subdifferential** of f at y and any element of $\partial f(y)$ is called a **subgradient** and denoted by $g_f(y)$.

(ii) We denote by $f_{\leq 0}$ the **sublevel set** of f , that is,

$$f_{\leq 0} := \{x \in \mathbb{R}^n \mid f(x) \leq 0\}. \quad (2.35)$$

Definition 2.16 Let $C \subseteq \mathbb{R}^n$. The **indicator function** of C at x is defined

$$I_C(x) := \begin{cases} 0 & \text{if } x \in C, \\ \infty & \text{otherwise.} \end{cases} \quad (2.36)$$

Definition 2.17 Let $C \subseteq \mathbb{R}^n$ be nonempty, closed and convex. We denote by $N_C(v)$ the **normal cone** of C at $v \in C$, i.e.,

$$N_C(v) := \{d \in \mathbb{R}^n \mid \langle d, y - v \rangle \leq 0 \text{ for all } y \in C\}. \quad (2.37)$$

Remark 2.18 It is well known that $\partial(I_C) = N_C$.

Example 2.19 Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function with a nonempty sublevel set $f_{\leq 0}$ (see (2.35)). Define an operator $\Pi_{f_{\leq 0}} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ by

$$\Pi_{f_{\leq 0}}(y) := \begin{cases} y - \frac{f(y)}{\|g_f(y)\|^2} g_f(y) & \text{if } f(y) > 0, \\ y & \text{if } f(y) \leq 0, \end{cases} \quad (2.38)$$

where $g_f(y)$ is a subgradient of f at y . The operator $\Pi_{f_{\leq 0}}$ is called the **subgradient projection** relative to f .

For a subgradient $g_f(y)$, we denote

$$L = L_f(y, g_f(y)) := \{x \in \mathbb{R}^n \mid f(y) + \langle g_f(y), x - y \rangle \leq 0\}. \quad (2.39)$$

Lemma 2.20 Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function and let $y \in \mathbb{R}^n$. Assume that $f_{\leq 0} \neq \emptyset$. Then the following assertions hold:

- (i) $f_{\leq 0} \subseteq L$. If $g_f(y) \neq 0$, then L is a half-space, otherwise $L = \mathbb{R}^n$.
- (ii) $\Pi_{f_{\leq 0}}(y) = P_L(y)$ consequently, $\Pi_{f_{\leq 0}}$ is a cutter and $\text{Fix}(\Pi_{f_{\leq 0}}) = f_{\leq 0}$.
- (iii) $I - \Pi_{f_{\leq 0}}$ is closed at 0.

Proof. See [4, Lemma 7.3], [15, Lemma 2.4] and [11, Lemma 4.2.5 and Corollary 4.2.6]. ■

The next class of operators was introduced by Aharoni et. al. in [1] for solving the convex feasibility problem. Later Gibali [24] and Censor and Gibali [14] used them for solving variational inequalities.

Let $C \subseteq \mathbb{R}^n$ be nonempty, closed and convex. Assume that C can be represented as a sublevel set of some convex function $c : \mathbb{R}^n \rightarrow \mathbb{R}$, that is,

$$C = \{x \in \mathbb{R}^n \mid c(x) \leq 0\}. \quad (2.40)$$

Given a point $z \in \mathbb{R}^n$ and a positive real number δ , we define for $z \notin C$ the ball

$$B(z, \delta c(z)) := \{x \in \mathbb{R}^n \mid \|x - z\| \leq \delta c(z)\}. \quad (2.41)$$

For $x, y \in \mathbb{R}^n$ we consider the set $H(x, y)$ as in (2.5) and define

$$A_\delta(z) := \{y \in \mathbb{R}^n \mid C \subseteq H(z, y) \text{ and } \text{int } B(z, \delta c(z)) \cap H(z, y) = \emptyset\}. \quad (2.42)$$

We also need to impose the following condition.

Condition 2.21 Given a set $C \subseteq \mathbb{R}^n$, described as in (2.40), we assume that for every $z \notin C$,

$$B(z, \delta c(z)) \cap C = \emptyset. \quad (2.43)$$

Every convex set C can be described by (2.40). We may take, for example, $c(z) = \text{dist}(z, C)$. In this case Condition 2.21 always holds for $\delta \in (0, 1)$.

Example 2.22 Given a nonempty, closed and convex subset $C \subseteq \mathbb{R}^n$, with the representation (2.40), and a real number $\delta \in (0, 1]$ such that Condition 2.21 holds, we define the operator $T_{C,\delta}$ at any $z \in \mathbb{R}^n$ by

$$T_{C,\delta}(z) := \begin{cases} P_{H(z,y)}(z) & \text{if } z \notin C, \\ z & \text{if } z \in C, \end{cases} \quad (2.44)$$

where $H(z, y)$ is built from any selection of y from $A_\delta(z)$, and call it a C - δ operator.

Observe that the subgradient projector $\Pi_{f_{\leq 0}}$ is a $T_{C,\delta}$ operator; see [15, Lemma 2.8]. The fact that any C - δ operator is a cutter operator follows from its definition. For the closedness of $T_{C,\delta} - I$ at 0, see, *e.g.*, [15, Lemma 2.7]. Alternatively, we show that, for any bounded sequence $\{x^k\}_{k=0}^\infty$, implication (2.33) is satisfied with $T := T_{C,\delta}$. Let $U := \Pi_{c \leq 0}$, where $c : \mathbb{R}^n \rightarrow \mathbb{R}$ is a convex function with $C := \{x \in \mathbb{R}^n \mid c(x) \leq 0\} \neq \emptyset$. Then, following Lemma 2.20, $I - U$ is closed at 0 and $\text{Fix}(U) = C$. Now, let $\{x^k\}_{k=0}^\infty$ be bounded and let $\lim_{k \rightarrow \infty} \|T_{C,\delta}(x^k) - x^k\| = 0$. Then, from the inequality $\|\delta c(x^k)\| \leq \|T_{C,\delta}(x^k) - x^k\|$, it follows that $\lim_{k \rightarrow \infty} c(x^k) = 0$. Consequently, $\lim_{k \rightarrow \infty} \text{dist}(x^k, C) = 0$, by the continuity of c . We claim that $\lim_{k \rightarrow \infty} \|U(x^k) - x^k\| = 0$. Indeed, by the definition of the subgradient g_c and the Cauchy–Schwarz inequality, we have

$$\begin{aligned} 0 \leq c(x^k) &= c(x^k) - c(P_C(x^k)) \leq \langle g_c(x^k), x^k - P_C(x^k) \rangle \\ &\leq \|g_c(x^k)\| \|x^k - P_C(x^k)\| = \|g_c(x^k)\| \text{dist}(x^k, C) \end{aligned} \quad (2.45)$$

and

$$\|U(x^k) - x^k\| = \frac{|c(x^k)|}{\|g_c(x^k)\|} \leq \text{dist}(x^k, C). \quad (2.46)$$

Consequently, $\lim_{k \rightarrow \infty} \|U(x^k) - x^k\| = 0$, as claimed.

3 The algorithm

Let $\mathcal{F} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a cutter. We need to assume that the following conditions hold in order to prove the convergence of our algorithm. These conditions were assumed to hold in [22] for solving $\text{VIP}(\mathcal{F}, S)$ (see (1.1)). Furthermore, the first two of these conditions guarantee that $\text{VIP}(\mathcal{F}, \text{Fix}(T))$ has a unique solution (see [20, Theorem 2.3.3] or [27, Chap. I, Corollary 4.3]).

Condition 3.1 $\mathcal{F}$ is continuous on $(\text{Fix}(T))_\varepsilon$ for some $\varepsilon > 0$.

Condition 3.2 $\mathcal{F}$ is α -strongly monotone on $(\text{Fix}(T))_\varepsilon$ for some $\varepsilon > 0$ and $\alpha > 0$, i.e.,

$$\langle \mathcal{F}(x) - \mathcal{F}(y), x - y \rangle \geq \alpha \|x - y\|^2 \text{ for all } x, y \in (\text{Fix}(T))_\varepsilon. \quad (3.1)$$

Condition 3.3 For some $q \in \text{Fix}(T)$, there exist some $\beta > 0$ and a bounded subset $E \subseteq \mathbb{R}^n$ such that

$$\langle \mathcal{F}(x), x - q \rangle \geq \beta \|\mathcal{F}(x)\| \text{ for all } x \notin E. \quad (3.2)$$

Condition 3.4 $I - T$ is closed at 0.

Remark 3.5 Conditions of the type of Condition 3.3 are commonly used in Optimization Theory (see, e.g., [21, Section 2.5] and [23, Section 8.3], where also examples of methods employing these conditions are presented). As it was observed by Fukushima in [22], a sufficient condition for Condition 3.3 to hold is that the vectors $\mathcal{F}(x)$ and x make an acute angle, which is uniformly bounded away from $\pi/2$, as $\|x\| \rightarrow \infty$. Indeed, Let $c \in (0, 1)$ and $r > 0$ be such that

$$\left\langle \frac{\mathcal{F}(x)}{\|\mathcal{F}(x)\|}, \frac{x}{\|x\|} \right\rangle \geq c \quad (3.3)$$

for $\|x\| \geq r$. Let $\beta \in (0, c)$, $q \in \text{Fix} T$ and $R \geq r$ be such that $\frac{\|q\|}{R} + \beta \leq c$. Then, for all $\|x\| \geq R$ we obtain

$$\begin{aligned} \langle \mathcal{F}(x), x - q \rangle &= \left(\left\langle \frac{\mathcal{F}(x)}{\|\mathcal{F}(x)\|}, \frac{x}{\|x\|} \right\rangle - \left\langle \frac{\mathcal{F}(x)}{\|\mathcal{F}(x)\|}, \frac{q}{\|x\|} \right\rangle \right) \|\mathcal{F}(x)\| \cdot \|x\| \\ &\geq \left(c - \frac{\|q\|}{R} \right) \|\mathcal{F}(x)\| \cdot \|x\| \geq \beta \|\mathcal{F}(x)\|. \end{aligned} \quad (3.4)$$

In addition, observe that Conditions 3.1 and 3.2 concern the behavior of $\mathcal{F}$ on $(\text{Fix}(T))_\varepsilon$, while Condition 3.3 deals with a rather global behavior.

Example 3.6 Let $\mathbb{R}^n$ be equipped with the standard inner product $\langle x, y \rangle := x^\top y$, $x, y \in \mathbb{R}^n$, $\lambda_1, \lambda_2 \in \mathbb{R}$ be such that $0 < \lambda_1 \leq \lambda_2$. Let $c \in (0, \lambda_1/\lambda_2)$, $a \in \mathbb{R}^n$ and $r := \frac{\lambda_2(1+c)\|a\|}{\lambda_1 - \lambda_2 c}$. Define $\mathcal{F} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ by

$$\mathcal{F}(x) = \begin{cases} \text{arbitrary} & \text{if } \|x\| \leq r \\ G(x)(x - a) & \text{if } \|x\| > r, \end{cases}$$

where $G(x)$ is a positive definite matrix with $\inf_{\|x\| > r} \lambda_{\min}(G(x)) \geq \lambda_1 > 0$ and $\sup_{\|x\| > r} \lambda_{\max}(G(x)) \leq \lambda_2$, where $\lambda_{\min}(G(x))$ and $\lambda_{\max}(G(x))$ denote the smallest and the largest eigenvalue of $G(x)$, respectively. We show, that $\mathcal{F}$ satisfies (3.3) for all $x \in \mathbb{R}^n$ with $\|x\| > r$. It follows from the inequalities

$$\|G(x)x\| \leq \|G(x)\| \cdot \|x\| = \lambda_{\max}(G(x))\|x\|$$

and

$$\lambda_{\min}(G(x))\|x\|^2 \leq x^\top G(x)x \leq \lambda_{\max}(G(x))\|x\|^2,$$

the Cauchy–Schwarz inequality, the triangle inequality and the monotonicity of the function $\xi \rightarrow \frac{\lambda_1 \xi - \lambda_2 \|a\|}{\lambda_2(\xi + \|a\|)}$ for $\xi > 0$, that

$$\begin{aligned} \frac{\langle \mathcal{F}(x), x \rangle}{\|\mathcal{F}(x)\| \cdot \|x\|} &= \frac{(x - a)^\top G(x)x}{\|G(x)(x - a)\| \cdot \|x\|} = \frac{x^\top G(x)x - a^\top G(x)x}{\|G(x)(x - a)\| \cdot \|x\|} \\ &\geq \frac{\lambda_{\min}(G(x))\|x\|^2 - \|a\|\lambda_{\max}(G(x))\|x\|}{\lambda_{\max}(G(x))(\|x\| + \|a\|)\|x\|} \\ &\geq \frac{\lambda_1\|x\| - \lambda_2\|a\|}{\lambda_2(\|x\| + \|a\|)} \geq c \end{aligned}$$

for all $x \in \mathbb{R}^n$ with $\|x\| > r$. If $G(x) = G$ for all $x \in \mathbb{R}^n$, then $\mathcal{F}(x) = G(x - a)$ and the unique solution of $\text{VIP}(\mathcal{F}, C)$ is $P_C^G a := \operatorname{argmin}_{x \in C} \|x - a\|_G$, where $\|\cdot\|_G$ denotes the norm induced by G , i.e., $\|u\|_G = (u^\top G u)^{\frac{1}{2}}$.

Let $\{\rho_k\}_{k=0}^\infty$ be a sequence of positive numbers satisfying

$$\lim_{k \rightarrow \infty} \rho_k = 0 \text{ and } \sum_{k=1}^\infty \rho_k = +\infty. \quad (3.5)$$

Algorithm 3.7

Initialization: Choose an arbitrary initial point $x^0 \in \mathbb{R}^n$ and set $k = 0$.

Iteration step: Given the current iterate x^k ,

(1) build the set $H_k := H(x^k, T(x^k))$ and calculate the “shifted point”

$$z^k := \begin{cases} x^k - \rho_k \mathcal{F}(x^k) / \|\mathcal{F}(x^k)\| & \text{if } \mathcal{F}(x^k) \neq 0, \\ x^k & \text{if } \mathcal{F}(x^k) = 0. \end{cases} \quad (3.6)$$

(2) Choose $\alpha_k \in [\mu, 2 - \mu]$ for some $\mu \in (0, 1)$ and calculate the next iterate as follows:

$$x^{k+1} = P_{\alpha_k}(z^k), \quad (3.7)$$

where $P_{\alpha_k} = I + \alpha_k(P_{H_k} - I)$ and P_{H_k} is the metric projection of $\mathbb{R}^n$ onto H_k .

(3) Set $k := k + 1$ and go to step (1).

Remark 3.8 Since T is a cutter, we have $\text{Fix}(T) \subseteq H_k$. Observe that (3.7) has an explicit form, because it is a relaxed projection onto a half-space ($x^k \neq T(x^k)$):

$$x^{k+1} = P_{\alpha_k}(z^k) = \begin{cases} z^k - \alpha_k \frac{\langle z^k - T(x^k), x^k - T(x^k) \rangle}{\|x^k - T(x^k)\|^2} (x^k - T(x^k)) & \text{if } z^k \notin H_k, \\ z^k & \text{if } z^k \in H_k. \end{cases} \quad (3.8)$$

An illustration of the iterative step of Algorithm 3.7 is given in Figure 2. It is clear that

$$\|z^k - x^k\| = \begin{cases} \rho_k & \text{if } \mathcal{F}(x^k) \neq 0, \\ 0 & \text{if } \mathcal{F}(x^k) = 0. \end{cases} \quad (3.9)$$

Therefore (3.5) yields $\lim_{k \rightarrow \infty} \|z^k - x^k\| = 0$.

4 Convergence

The following lemma is a consequence of Theorem 2.4, where P_{α_k} is a relaxation of P_{H_k} , both defined in Algorithm 3.7. Nevertheless, below we present a complete proof of this lemma for the convenience of the reader.

Lemma 4.1 *Let $y \in \mathbb{R}^n$ be arbitrary and let $\alpha_k \in (0, 2)$. Then in the setting of Algorithm 3.7 we have*

$$\|P_{\alpha_k}(y) - w\|^2 \leq \|y - w\|^2 - \frac{2 - \alpha_k}{\alpha_k} \|P_{\alpha_k}(y) - y\|^2 \text{ for all } w \in \text{Fix}(T). \quad (4.1)$$




Figure 2: Iterative step of Algorithm 3.7

Consequently,

$$\text{dist}(P_{\alpha_k}(y), \text{Fix}(T))^2 \leq \text{dist}(y, \text{Fix}(T))^2 - \frac{2 - \alpha_k}{\alpha_k} \|P_{\alpha_k}(y) - y\|^2. \quad (4.2)$$

Proof. Let $w \in \text{Fix}(T)$. Since $\text{Fix}(T) \subseteq H_k$, the characterization of the metric projection yields

$$\langle y - P_{H_k}(y), y - w \rangle \geq \|P_{H_k}(y) - y\|^2 = \frac{1}{\alpha_k^2} \|P_{\alpha_k}(y) - y\|^2 \quad (4.3)$$

and therefore we have

$$\begin{aligned} \|P_{\alpha_k}(y) - w\|^2 &= \|y + \alpha_k(P_{H_k}(y) - y) - w\|^2 \\ &= \|y - w\|^2 + \alpha_k^2 \|P_{H_k}(y) - y\|^2 - 2\alpha_k \langle y - P_{H_k}(y), y - w \rangle \\ &\leq \|y - w\|^2 - \frac{2 - \alpha_k}{\alpha_k} \|P_{\alpha_k}(y) - y\|^2. \end{aligned} \quad (4.4)$$

When we set $w = P_{\text{Fix}(T)}(y)$ in (4.1), we obtain

$$\text{dist}(P_{\alpha_k}(y), \text{Fix}(T))^2 \leq \text{dist}(y, \text{Fix}(T))^2 - \frac{2 - \alpha_k}{\alpha_k} \|P_{\alpha_k}(y) - y\|^2 \quad (4.5)$$

and the proof is complete. ■

Lemma 4.2 *Assume that Condition 3.3 holds. Then any sequence $\{x^k\}_{k=0}^\infty$ generated by Algorithm 3.7 is bounded.*

Proof. The proof is structured along the lines of [22, Lemma 3]. Let $q \in \text{Fix}(T)$, $\beta > 0$ and a bounded subset $E \subseteq \mathbb{R}^n$ be such that (3.2) is satisfied. We show that, for sufficiently large k , we have

$$\|x^{k+1} - q\| \leq \|x^k - q\| \quad (4.6)$$

if $x^k \notin E$ and

$$\|x^{k+1} - q\| \leq \|x^k - q\| + \rho \quad (4.7)$$

otherwise, where $\rho = \sup_k \rho_k$. This implies that $\{x^k\}_{k=0}^\infty$ is bounded. If $\mathcal{F}(x^k) = 0$, then $x^k = z^k$. By the definition of x^{k+1} and Lemma 4.1, applied to $y := z^k$, $w := q$, inequality (4.6) is satisfied. Now assume that $\mathcal{F}(x^k) \neq 0$. Lemma 4.1 implies that, for each $z^k \in \mathbb{R}^n$,

$$\|P_{\alpha_k}(z^k) - q\|^2 \leq \|z^k - q\|^2. \quad (4.8)$$

Therefore

$$\begin{aligned} \|x^{k+1} - q\|^2 &\leq \left\| x^k - \rho_k \frac{\mathcal{F}(x^k)}{\|\mathcal{F}(x^k)\|} - q \right\|^2 = \|x^k - q\|^2 \\ &\quad - 2 \frac{\rho_k}{\|\mathcal{F}(x^k)\|} \langle \mathcal{F}(x^k), x^k - q \rangle + \rho_k^2. \end{aligned} \quad (4.9)$$

Thus, if $x^k \notin E$, we have, by (4.9) and Condition 3.3,

$$\|x^{k+1} - q\|^2 \leq \|x^k - q\|^2 - 2\rho_k\beta + \rho_k^2 = \|x^k - q\|^2 - \rho_k(2\beta - \rho_k). \quad (4.10)$$

Since $\rho_k > 0$ and $\lim_{k \rightarrow \infty} \rho_k = 0$, the latter inequality implies (4.6), provided that k is sufficiently large. On the other hand, by (4.8), the definition of z^k (see (3.6)) and the triangle inequality, we obtain

$$\|x^{k+1} - q\| \leq \left\| (x^k - q) - \rho_k \frac{\mathcal{F}(x^k)}{\|\mathcal{F}(x^k)\|} \right\| \leq \|x^k - q\| + \rho_k, \quad (4.11)$$

i.e., (4.7) is satisfied. Hence $\{x^k\}_{k=0}^\infty$ is indeed bounded as we have already observed. ■

Lemma 4.3 *Assume that Condition 3.3 holds. Then any sequence $\{x^k\}_{k=0}^\infty$ generated by Algorithm 3.7 satisfies*

$$\lim_{k \rightarrow \infty} \text{dist}(x^k, \text{Fix}(T)) = 0. \quad (4.12)$$

Proof. Recall that $R := P_{\text{Fix}(T)}$. Lemma 4.1 implies that

$$\text{dist}(x^{k+1}, \text{Fix}(T)) \leq \text{dist}(z^k, \text{Fix}(T)). \quad (4.13)$$

If $\mathcal{F}(x^k) \neq 0$, then, by the triangle inequality,

$$\begin{aligned} \text{dist}(z^k, \text{Fix}(T)) &= \|z^k - R(x^k)\| = \left\| x^k - \rho_k \frac{\mathcal{F}(x^k)}{\|\mathcal{F}(x^k)\|} - R(x^k) \right\| \\ &\leq \|x^k - R(x^k)\| + \rho_k \\ &= \text{dist}(x^k, \text{Fix}(T)) + \rho_k. \end{aligned} \quad (4.14)$$

If $\mathcal{F}(x^k) = 0$, then $x^k = z^k$ and, consequently,

$$\text{dist}(z^k, \text{Fix}(T)) = \text{dist}(x^k, \text{Fix}(T)). \quad (4.15)$$

Therefore in both cases we have

$$\text{dist}(x^{k+1}, \text{Fix}(T)) \leq \text{dist}(x^k, \text{Fix}(T)) + \tilde{\rho}_k, \quad (4.16)$$

where

$$\tilde{\rho}_k := \begin{cases} \rho_k, & \text{if } \mathcal{F}(x^k) \neq 0, \\ 0, & \text{if } \mathcal{F}(x^k) = 0. \end{cases} \quad (4.17)$$

Define $a_k := \text{dist}(x^k, \text{Fix}(T))$. Let $C \subseteq \mathbb{R}^n$ be a bounded, closed and convex set such that $\{x^k\}_{k=0}^\infty \subseteq C$ and $\text{Fix}(T) \cap C \neq \emptyset$. The existence of such a set C follows from Lemma 4.2. By Proposition 2.6(iii) and Lemma 4.1,

$$D^2(a_k) = D^2(\text{dist}(x^k, \text{Fix}(T))) \leq \|x^k - T(x^k)\|^2 = \frac{1}{\alpha_k^2} \|x^k - P_{\alpha_k}(x^k)\|^2 \quad (4.18)$$

$$\leq \frac{1}{\alpha_k(2 - \alpha_k)} \left((\text{dist}(x^k, \text{Fix}(T)))^2 - (\text{dist}(P_{\alpha_k}(x^k), \text{Fix}(T)))^2 \right). \quad (4.19)$$

On the other hand, by the nonexpansivity of P_{α_k} , (3.9) and (4.17), we get

$$\|x^{k+1} - P_{\alpha_k}(x^k)\|^2 = \|P_{\alpha_k}(z^k) - P_{\alpha_k}(x^k)\|^2 \leq \|z^k - x^k\|^2 = \tilde{\rho}_k^2. \quad (4.20)$$

Therefore

$$\|x^{k+1} - P_{\alpha_k}(x^k)\| \leq \tilde{\rho}_k. \quad (4.21)$$

Let $s^k = R(P_{\alpha_k}(x^k))$, i.e.,

$$\|P_{\alpha_k}(x^k) - s^k\| = \text{dist}(P_{\alpha_k}(x^k), \text{Fix}(T)). \quad (4.22)$$

Then, by the triangle inequality, we get

$$\|x^{k+1} - s^k\| \leq \|x^{k+1} - P_{\alpha_k}(x^k)\| + \|P_{\alpha_k}(x^k) - s^k\|. \quad (4.23)$$

On the other hand, since $s^k \in \text{Fix}(T)$, we have

$$\text{dist}(x^{k+1}, \text{Fix}(T)) \leq \|x^{k+1} - s^k\|. \quad (4.24)$$

From the last four inequalities we get

$$\begin{aligned} a_{k+1} &\leq \|x^{k+1} - P_{\alpha_k}(x^k)\| + \text{dist}(P_{\alpha_k}(x^k), \text{Fix}(T)) \\ &\leq \tilde{\rho}_k + \text{dist}(P_{\alpha_k}(x^k), \text{Fix}(T)) \end{aligned} \quad (4.25)$$

or, equivalently,

$$(\text{dist}(P_{\alpha_k}(x^k), \text{Fix}(T)))^2 \geq a_{k+1}^2 - \tilde{\rho}_k^2 - 2\tilde{\rho}_k \text{dist}(P_{\alpha_k}(x^k), \text{Fix}(T)). \quad (4.26)$$

Using the above inequality for (4.19), we get for all $k \geq 0$,

$$D^2(a_k) \leq \frac{1}{\alpha_k(2 - \alpha_k)} (a_k^2 - a_{k+1}^2 + \tilde{\rho}_k^2 + 2\tilde{\rho}_k \text{dist}(P_{\alpha_k}(x^k), \text{Fix}(T))). \quad (4.27)$$

Now, by Lemma 4.2, the sequence $\{x^k\}_{k=0}^\infty$ is bounded and, therefore, so is $\{a_k\}_{k=0}^\infty$. By Lemma 4.1 with $y := x^k$, we have

$$\begin{aligned} &(\text{dist}(P_{\alpha_k}(x^k), \text{Fix}(T)))^2 \\ &\leq \|P_{\alpha_k}(x^k) - z\|^2 \leq \|x^k - z\|^2 - \frac{2 - \alpha_k}{\alpha_k} \|P_{\alpha_k}(x^k) - x^k\|^2 \leq \|x^k - z\|^2 \end{aligned} \quad (4.28)$$

for all $z \in \text{Fix}(T)$. Taking $z := R(x^k)$ in the above inequalities, we obtain

$$\text{dist}(P_{\alpha_k}(x^k), \text{Fix}(T)) \leq a_k = \text{dist}(x^k, \text{Fix}(T)). \quad (4.29)$$

Therefore the sequence $\{\text{dist}(P_{\alpha_k}(x^k), \text{Fix}(T))\}_{k=0}^\infty$ is also bounded. Since $\alpha_k \in [\mu, 2 - \mu]$ for some $\mu \in (0, 1)$, we have $1/(\alpha_k(2 - \alpha_k)) \leq 1/\mu^2$. Denote $b_k := \tilde{\rho}_k^2 + 2\tilde{\rho}_k \text{dist}(P_{\alpha_k}(x^k), \text{Fix}(T))$. Using (4.27), we get

$$\begin{aligned} D^2(a_k) &\leq \frac{1}{\mu^2} (a_k^2 - a_{k+1}^2 + b_k) \\ &= \frac{1}{\mu^2} (a_k + a_{k+1})(a_k - a_{k+1}) + \frac{1}{\mu^2} b_k. \end{aligned} \quad (4.30)$$

Since $\{a_k\}_{k=0}^\infty$ is bounded, we have $1/\mu^2 (a_k + a_{k+1}) \leq M$ for some $M > 0$. By the definition of $\tilde{\rho}_k$, $\lim_{k \rightarrow \infty} b_k = 0$. Hence

$$\frac{1}{M} D^2(a_k) \leq a_k - a_{k+1} + \frac{1}{\mu^2 M} b_k \quad (4.31)$$

and now we can apply Lemma 2.13 to conclude that $\lim_{k \rightarrow \infty} \text{dist}(x^k, \text{Fix}(T)) = 0$, which completes the proof. ■

Lemma 4.4 *Assume that Condition 3.3 holds. Any sequence $\{x^k\}_{k=0}^\infty$ generated by Algorithm 3.7 satisfies*

$$\lim_{k \rightarrow \infty} \|x^{k+1} - x^k\| = 0. \quad (4.32)$$

Proof. If $\mathcal{F}(x^k) \neq 0$, then by the triangle inequality, the nonexpansivity of P_{α_k} and by (3.9), we obtain for all $k \geq 0$,

$$\begin{aligned} \|x^{k+1} - x^k\| &\leq \|x^{k+1} - P_{\alpha_k}(x^k)\| + \|P_{\alpha_k}(x^k) - x^k\| \\ &= \|P_{\alpha_k}(z^k) - P_{\alpha_k}(x^k)\| + \|P_{\alpha_k}(x^k) - x^k\| \\ &\leq \|z^k - x^k\| + \|P_{\alpha_k}(x^k) - x^k\| \\ &\leq \rho_k + \alpha_k \text{dist}(x^k, H_k), \end{aligned} \quad (4.33)$$

where the latter inequality follows from (3.9) and from the obvious equality $\alpha_k \text{dist}(x^k, H_k) = \|P_{\alpha_k}(x^k) - x^k\|$. Since for all $k \geq 0$, $\text{Fix}(T) \subseteq H_k$, we have

$$\text{dist}(x^k, H_k) \leq \text{dist}(x^k, \text{Fix}(T)). \quad (4.34)$$

Thus,

$$\|x^{k+1} - x^k\| \leq \rho_k + \alpha_k \operatorname{dist}(x^k, \operatorname{Fix}(T)). \quad (4.35)$$

In case $\mathcal{F}(x^k) = 0$, we have

$$\|x^{k+1} - x^k\| = \alpha_k \operatorname{dist}(x^k, H_k) \leq \alpha_k \operatorname{dist}(x^k, \operatorname{Fix}(T)). \quad (4.36)$$

By Lemma 4.3 and (3.5), we obtain the required result. ■

Theorem 4.5 *Assume that Conditions 3.1 – 3.4 are satisfied. Then any sequence $\{x^k\}_{k=0}^\infty$ generated by Algorithm 3.7 converges to the unique solution x^* of $\operatorname{VIP}(\mathcal{F}, \operatorname{Fix}(T))$.*

Proof. Let x^* be the unique solution of problem (1.3). By Lemma 4.3, $x^k \in (\operatorname{Fix}(T))_\varepsilon$ for all sufficiently large k , where $(\operatorname{Fix}(T))_\varepsilon$ is the set given in Conditions 3.1 and 3.2 (without loss of generality, ε is the same in both conditions). By Condition 3.2, we have

$$\langle \mathcal{F}(x^k) - \mathcal{F}(x^*), x^k - x^* \rangle \geq \alpha \|x^k - x^*\|^2 \quad (4.37)$$

and

$$\begin{aligned} \langle \mathcal{F}(x^k) - \mathcal{F}(x^*), x^k - x^* \rangle &= \langle \mathcal{F}(x^k), x^k - x^{k+1} \rangle + \langle \mathcal{F}(x^k), x^{k+1} - x^* \rangle \\ &\quad - \langle \mathcal{F}(x^*), x^k - x^* \rangle. \end{aligned} \quad (4.38)$$

Therefore

$$\langle \mathcal{F}(x^k), x^{k+1} - x^* \rangle \geq \alpha \|x^k - x^*\|^2 + \langle \mathcal{F}(x^*), x^k - x^* \rangle + \langle \mathcal{F}(x^k), x^{k+1} - x^* \rangle. \quad (4.39)$$

Let λ be an arbitrary positive number. Then we have

$$\langle \mathcal{F}(x^*), x^k - x^* \rangle \geq -\lambda \quad (4.40)$$

for all sufficiently large k . Indeed. The inequality is clear if $\mathcal{F}(x^*) = 0$. Assume now that $\mathcal{F}(x^*) \neq 0$ and let $\varepsilon := \lambda / \|\mathcal{F}(x^*)\|$. By Lemma 4.3, we have $\|x^k - z^k\| \leq \varepsilon$ for all sufficiently large k , where $z^k = P_{\operatorname{Fix}(T)}(x^k)$. Consequently,

$$\begin{aligned} \langle \mathcal{F}(x^*), x^k - x^* \rangle &= \langle \mathcal{F}(x^*), z^k - x^* \rangle + \langle \mathcal{F}(x^*), x^k - z^k \rangle \\ &\geq -\|\mathcal{F}(x^*)\| \cdot \|x^k - z^k\| \geq -\|\mathcal{F}(x^*)\| \cdot \varepsilon = -\lambda. \end{aligned}$$

By the Cauchy–Schwarz inequality,

$$\langle \mathcal{F}(x^k), x^{k+1} - x^k \rangle \geq -\|\mathcal{F}(x^k)\| \|x^{k+1} - x^k\|. \quad (4.41)$$

From the boundedness of $\{x^k\}_{k=0}^\infty$ (see Lemma 4.2) and from Lemma 4.3, it follows, due to the continuity of $\mathcal{F}$ on $(\text{Fix}(T))_\varepsilon$ (Condition 3.1), that the sequence $\{\mathcal{F}(x^k)\}_{k=0}^\infty$ is also bounded. Lemma 4.4 and inequality (4.41) guarantee that

$$\langle \mathcal{F}(x^k), x^{k+1} - x^k \rangle \geq -\lambda \quad (4.42)$$

for all sufficiently large k . Applying (4.40) and (4.42) to (4.39), we obtain

$$\langle \mathcal{F}(x^k), x^{k+1} - x^* \rangle \geq \alpha \|x^k - x^*\|^2 - 2\lambda \quad (4.43)$$

for all sufficiently large k . Divide the indices of $\{x^k\}_{k=0}^\infty$ as follows:

$$\Gamma := \{k \geq 0 \mid \mathcal{F}(x^k) = 0\} \text{ and } \tilde{\Gamma} := \{k \geq 0 \mid \mathcal{F}(x^k) \neq 0\}. \quad (4.44)$$

Equation (4.43) implies, since λ is arbitrary, that for $k \in \Gamma$,

$$\lim_{k \rightarrow \infty} x^k = x^*. \quad (4.45)$$

We now show that the sequence $\{x^k\}_{k \in \tilde{\Gamma}}$ contains a subsequence which converges to x^* . To this end, let us consider the indices in $\tilde{\Gamma}$ and suppose that there exist $\zeta > 0$ and an integer k_0 such that

$$\|x^k - x^*\| \geq \zeta \text{ for all } k \in \tilde{\Gamma}, k \geq k_0. \quad (4.46)$$

By Lemma 4.1,

$$\begin{aligned} \|x^{k+1} - x^*\|^2 &= \|P_{\alpha_k}(z^k) - x^*\|^2 \leq \|z^k - x^*\|^2 - \frac{2 - \alpha_k}{\alpha_k} \|P_{\alpha_k}(z^k) - z^k\|^2 \\ &= \left\| \left(x^k - \rho_k \frac{\mathcal{F}(x^k)}{\|\mathcal{F}(x^k)\|} \right) - x^* \right\|^2 \\ &\quad - \frac{2 - \alpha_k}{\alpha_k} \left\| x^{k+1} - \left(x^k - \rho_k \frac{\mathcal{F}(x^k)}{\|\mathcal{F}(x^k)\|} \right) \right\|^2 \\ &= \|x^k - x^*\|^2 - \frac{2 - \alpha_k}{\alpha_k} \|x^{k+1} - x^k\|^2 - 2 \frac{\rho_k}{\|\mathcal{F}(x^k)\|} \langle \mathcal{F}(x^k), x^k - x^* \rangle \\ &\quad + 2 \frac{(2 - \alpha_k) \rho_k}{\alpha_k \|\mathcal{F}(x^k)\|} \langle \mathcal{F}(x^k), x^{k+1} - x^k \rangle + \rho_k^2 \left(1 - \frac{2 - \alpha_k}{\alpha_k} \right). \end{aligned} \quad (4.47)$$

This is less than or equal to

$$\begin{aligned}
& \|x^k - x^*\|^2 - 2 \frac{\rho_k}{\|\mathcal{F}(x^k)\|} \langle \mathcal{F}(x^k), x^k - x^* \rangle \\
& + 2 \frac{(2 - \alpha_k) \rho_k}{\alpha_k \|\mathcal{F}(x^k)\|} \langle \mathcal{F}(x^k), x^{k+1} - x^k \rangle + \rho_k^2 \left(1 - \frac{2 - \alpha_k}{\alpha_k} \right) \\
& = \|x^k - x^*\|^2 - 2 \frac{\rho_k}{\|\mathcal{F}(x^k)\|} \langle \mathcal{F}(x^k), x^{k+1} - x^* \rangle \\
& + 2 \left(\frac{2 - \alpha_k}{\alpha_k} + 1 \right) \frac{\rho_k}{\|\mathcal{F}(x^k)\|} \langle \mathcal{F}(x^k), x^{k+1} - x^k \rangle \\
& + \rho_k^2 \left(1 - \frac{2 - \alpha_k}{\alpha_k} \right). \tag{4.48}
\end{aligned}$$

So, combining the above relations, we obtain

$$\begin{aligned}
\|x^{k+1} - x^*\|^2 & \leq \|x^k - x^*\|^2 - 2 \frac{\rho_k}{\|\mathcal{F}(x^k)\|} \langle \mathcal{F}(x^k), x^{k+1} - x^* \rangle \\
& + 2 \left(\frac{2 - \alpha_k}{\alpha_k} + 1 \right) \frac{\rho_k}{\|\mathcal{F}(x^k)\|} \langle \mathcal{F}(x^k), x^{k+1} - x^k \rangle \\
& + \rho_k^2 \left(1 - \frac{2 - \alpha_k}{\alpha_k} \right). \tag{4.49}
\end{aligned}$$

Since $\{\mathcal{F}(x^k)\}_{k=0}^\infty$ is bounded, there exists $\tau > 0$ such that $\|\mathcal{F}(x^k)\| \leq \tau$. Therefore,

$$-\frac{1}{\|\mathcal{F}(x^k)\|} \leq -\frac{1}{\tau}, \text{ for all } k \in \tilde{\Gamma}. \tag{4.50}$$

Since λ is arbitrary, we can assume that

$$2\lambda \leq \frac{1}{4} \alpha \zeta^2. \tag{4.51}$$

By similar arguments as in derivation of inequality (4.42) and by the boundedness of the sequence $\{(2 - \alpha_k) / \alpha_k\}_{k=0}^\infty$, we can assume that, for all sufficiently large k ,

$$\left(\frac{2 - \alpha_k}{\alpha_k} + 1 \right) \langle \mathcal{F}(x^k), x^{k+1} - x^k \rangle \leq \frac{1}{4} \alpha \zeta^2 - 2\lambda. \tag{4.52}$$

Since $\rho_k \rightarrow 0$ and again using the boundedness of $(2 - \alpha_k) / \alpha_k$, we can also assume that, for all sufficiently large k ,

$$\rho_k \left(1 - \frac{2 - \alpha_k}{\alpha_k} \right) \leq \frac{\alpha \zeta^2}{2\tau}. \tag{4.53}$$

Applying (4.43), (4.52) and (4.53) to (4.49), we get

$$\begin{aligned}
\|x^{k+1} - x^*\|^2 &\leq \|x^k - x^*\|^2 - 2 \frac{\rho_k}{\|\mathcal{F}(x^k)\|} (\alpha\zeta^2 - 2\lambda) \\
&\quad + 2 \frac{\rho_k}{\|\mathcal{F}(x^k)\|} \left(\frac{1}{4} \alpha\zeta^2 - 2\lambda \right) + \rho_k \frac{\alpha\zeta^2}{2\tau} \\
&= \|x^k - x^*\|^2 - \frac{3}{2} \frac{\rho_k}{\|\mathcal{F}(x^k)\|} \alpha\zeta^2 + \rho_k \frac{\alpha\zeta^2}{2\tau}.
\end{aligned} \tag{4.54}$$

Combining (4.50) with (4.54), we get

$$\|x^{k+1} - x^*\|^2 \leq \|x^k - x^*\|^2 - \rho_k \frac{\alpha\zeta^2}{\tau} \tag{4.55}$$

for all sufficiently large k . Thus there exists an integer $\tilde{k} \in \tilde{\Gamma}$ such that

$$\|x^{k+1} - x^*\|^2 \leq \|x^k - x^*\|^2 - \rho_k \frac{\alpha\zeta^2}{\tau} \text{ for all } k \in \tilde{\Gamma} \text{ such that } k \geq \tilde{k}. \tag{4.56}$$

By adding these inequalities for $k = \tilde{k}, \tilde{k} + 1, \dots, \tilde{k} + \ell$ over $k \in \tilde{\Gamma}$, we obtain

$$\|x^{\tilde{k}+\ell+1} - x^*\|^2 \leq \|x^{\tilde{k}} - x^*\|^2 - \frac{\alpha\zeta^2}{\tau} \sum_{k \in \tilde{\Gamma}, k=\tilde{k}}^{\tilde{k}+\ell} \rho_k \tag{4.57}$$

for any $\ell > 0$. However, this is impossible in view of (3.5). So there exists no $\zeta > 0$ such that (4.46) is satisfied. Therefore $\{x^k\}_{k \in \tilde{\Gamma}}$ contains a subsequence $\{x^k\}_{k \in \hat{\Gamma}}$, $\hat{\Gamma} \subseteq \tilde{\Gamma}$, converging to x^* , i.e., there is a subsequence $\{x^k\}_{k \in \Gamma \cup \hat{\Gamma}}$ of the whole sequence $\{x^k\}_{k=0}^\infty$ which converges to x^* . In order to prove that the entire sequence $\{x^k\}_{k=0}^\infty$ converges to x^* , suppose to the contrary that there exists a subsequence of $\{x^k\}_{k=0}^\infty$ converging to $\hat{x}$ and $\hat{x} \neq x^*$. By Lemma 4.4, $\lim_{k \rightarrow \infty} \|x^{k+1} - x^k\| = 0$, therefore, there exists $\zeta > 0$ and an arbitrarily large integer $j \in \tilde{\Gamma}$ such that

$$\|x^j - x^*\| \geq \zeta \text{ and } \|x^{j+1} - x^*\| \geq \|x^j - x^*\|. \tag{4.58}$$

However, if j is sufficiently large, we may apply an argument similar to that used to derive (4.56) and obtain the inequality

$$\|x^{j+1} - x^*\| < \|x^j - x^*\|, \tag{4.59}$$

which contradicts (4.58). Therefore the whole sequence $\{x^k\}_{k=0}^\infty$ does converge to x^* , as asserted. ■

Remark 4.6 In [34] the operator $\mathcal{F}$ is assumed to be Lipschitz continuous and strongly monotone on the image of T , while here $\mathcal{F}$ is only assumed to be continuous on $(\text{Fix}(T))_\varepsilon$ for some $\varepsilon > 0$. In [34, Theorem 5] Yamada and Ogura showed that the strong monotonicity of $\mathcal{F}$ could be weakened and replaced by the paramonotonicity. In [35] Yamada et al. applied successfully Algorithm (1.4) to the minimization of the Moreau envelope of nonsmooth convex functions, where only Lipschitz continuity and paramonotonicity were assumed.

5 An application

Given an operator $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, we would like to find its minimizers. Clearly, we cannot look for an optimal solution as defined for a scalar optimization problem ($m = 1$). Therefore we need to define *a priori* which solution concept is chosen. One might consider the lexicographic order, denoted by $\preceq_L$. This partial order is defined for $x, y \in \mathbb{R}^m$ as follows:

$$x \preceq_L y \Leftrightarrow x = y \text{ or } x_k < y_k \text{ where } k := \min\{i = 1, \dots, m \mid x_i \neq y_i\}. \quad (5.1)$$

Now consider the case where $m = 2$, i.e., $f : \mathbb{R}^n \rightarrow \mathbb{R}^2$, and denote by $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$ the i -th coordinate ($i = 1, 2$) of the function f . Then our goal is to minimize f with respect to $\preceq_L$. This problem is also called a *two-stage* or a *bi-level* or a *hierarchical optimization problem*. Before introducing the connection of this problem to our VIP, we recall some definitions and properties.

Definition 5.1 Let $A : \mathbb{R}^n \rightarrow 2^{\mathbb{R}^n}$ be a set-valued mapping.

(i) A is called a **maximal monotone mapping** if it is monotone, i.e.,

$$\langle u - v, x - y \rangle \geq 0 \text{ for all } u \in A(x) \text{ and } v \in A(y), \quad (5.2)$$

and the **graph** $G(A)$ of A ,

$$G(A) := \{(x, u) \in \mathbb{R}^n \times \mathbb{R}^n \mid u \in A(x)\}, \quad (5.3)$$

is not properly contained in the graph of any other monotone mapping.

(ii) The **resolvent** of A with parameter λ is the operator $J_\lambda^A := (I + \lambda A)^{-1}$, where I is the identity operator.

Remark 5.2 It is well known that for $\lambda > 0$,

(i) A is monotone if and only if the resolvent J_λ^A of A is single-valued and firmly nonexpansive.

(ii) A is maximal monotone if and only if J_λ^A is single-valued, firmly nonexpansive and its domain is $\mathbb{R}^n$, where

$$\text{dom}(J_\lambda^A) := \{x \in \mathbb{R}^n \mid J_\lambda^A(x) \neq \emptyset\}. \quad (5.4)$$

(iii)

$$0 \in A(x) \Leftrightarrow x \in \text{Fix}(J_\lambda^A). \quad (5.5)$$

Example 5.3 Let $C \subseteq \mathbb{R}^n$ be nonempty, closed and convex. The metric projection onto C is precisely the resolvent of the normal cone mapping, *i.e.*,

$$P_C = J_\lambda^{N_C}. \quad (5.6)$$

In addition, it is known that N_C is a maximal monotone mapping.

Remark 5.4 Let $C \subseteq \mathbb{R}^n$ be closed and convex. If a function $g : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex, then

$$x^* \in \text{Argmin}\{g \mid C\} \iff 0 \in \partial g(x^*) + N_C(x^*), \quad (5.7)$$

(see [25, Chapter VII, Theorem 1.1.1]). In the case where g is continuously differentiable, it follows from the first order optimality condition that

$$x^* \in \text{Argmin}\{g \mid C\} \iff x^* \text{ solves } \text{VIP}(\nabla g, C), \quad (5.8)$$

see, *e.g.*, [7, Proposition 3.1, p. 210].

Remark 5.5 Both set-valued mappings, ∂g and N_C , are maximal monotone and $\text{dom}(\partial g) = \mathbb{R}^n$, hence also $\partial g + N_C$ is maximal monotone (see [6, Corollary 24.4 (i)]) and therefore

$$0 \in (\partial g + N_C)(x^*) \Leftrightarrow x^* \in \text{Fix}(J_\lambda^{\partial g + N_C}). \quad (5.9)$$

Let's go back to the hierarchical optimization problem:

$$\min\{f_2 \mid \text{Argmin}\{f_1 \mid C\}\}. \quad (5.10)$$

Under the assumption of the convexity of $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$, $i = 1, 2$, and the continuous differentiability of f_2 , problem (5.10) can be reformulated as

$\text{VIP}(\nabla f_2, \text{Fix}(J_\lambda^{\partial f_1 + N_C}))$. That is, we look for a point $x^* \in \text{Fix}(J_\lambda^{\partial f_1 + N_C})$ such that

$$\langle \nabla f_2(x^*), x - x^* \rangle \geq 0 \text{ for all } x \in \text{Fix}(J_\lambda^{\partial f_1 + N_C}). \quad (5.11)$$

So, under appropriate assumptions on $f : \mathbb{R}^n \rightarrow \mathbb{R}^2$, which assure that ∇f_2 satisfies Conditions 3.1 – 3.4, we could apply Algorithm 3.7.

Next we present an example that can be translated into an appropriate VIP over the fixed point set of a cutter operator.

Example 5.6 Let $C \subseteq \mathbb{R}^n$ be closed and convex. Given a convex function $g : \mathbb{R}^n \rightarrow \mathbb{R}$, we are interested in minimizing g over C so that the solution has minimal p -th norm, where $p \geq 2$. This solution is called a *p -minimal-norm solution*.

Define the operator $f = (f_1, f_2) : \mathbb{R}^n \rightarrow \mathbb{R}^2$ by

$$f = \left(\begin{array}{c} g \\ \frac{1}{p} \|\cdot\|_p^p + \frac{\alpha}{2} \|\cdot\|_2^2 \end{array} \right), \quad (5.12)$$

where $\|\cdot\|_p$ denotes the p -th norm, i.e., $\|x\|_p := (\sum_{i=1}^n |x_i|^p)^{1/p}$. Consider the following special case of problem (5.10):

$$\begin{cases} \text{minimize } \frac{1}{p} \|x\|_p^p + \frac{\alpha}{2} \|x\|_2^2 \\ \text{s.t. } x \in \text{Argmin}\{g(x) \mid C\}, \end{cases} \quad (5.13)$$

which is a regularization of the problem under consideration. Notice that f_2 is a sum of convex and strongly convex functions and, therefore, ∇f_2 is strongly monotone.

For $p = 2$ we get that $\nabla(\frac{1}{2} \|\cdot\|_2^2) = I$. Hence ∇f_2 is Lipschitz continuous on $\mathbb{R}^n$. Moreover, we do not need a regularization term to obtain strong monotonicity and we can set $\alpha = 0$. Therefore we can use Yamada's and Ogura's hybrid steepest descent algorithm (see [34, Section 4]) to solve $\text{VIP}(I, \text{Fix}(J_\lambda^{\partial g + N_C}))$ and obtain a 2-minimal-norm solution.

Let now $p > 2$. In this case $\nabla(\frac{1}{p} \|x\|_p^p) = (x_1|x_1|^{p-2}, x_2|x_2|^{p-2}, \dots, x_n|x_n|^{p-2})$. One can easily check that $\nabla(\frac{1}{p} \|\cdot\|_p^p)$ is not globally Lipschitz continuous. Therefore we cannot use Yamada's and Ogura's algorithm. However, we can use Algorithm 3.7 to solve $\text{VIP}(\nabla(\frac{1}{p} \|\cdot\|_p^p + \frac{\alpha}{2} \|\cdot\|_2^2), \text{Fix}(J_\alpha^{\partial g + N_C}))$. To see this, denote $\mathcal{F}_1 := \nabla(\frac{1}{p} \|\cdot\|_p^p)$, $\mathcal{F}_2 := \nabla(\frac{\alpha}{2} \|\cdot\|_2^2)$ and $\mathcal{F} := \mathcal{F}_1 + \mathcal{F}_2$. By Remark 3.5, it suffices to show that $\langle \mathcal{F}_i(x), x \rangle \geq c \|\mathcal{F}_i(x)\|_2 \|x\|_2$, for all $\|x\|_2 \geq R > 0$,

$i = 1, 2$, and some $c \in (0, 1)$. Notice that for $i = 2$ this inequality holds for all $c \in (0, 1)$, because

$$\langle \mathcal{F}_2(x), x \rangle = \alpha \|x\|_2^2 = \|\mathcal{F}_2(x)\|_2 \cdot \|x\|_2. \quad (5.14)$$

For $i = 1$ the inequality is equivalent to

$$(\|x\|_p^p)^2 \geq c \cdot \|x\|_{2p-2}^{2p-2} \cdot \|x\|_2^2, \quad (5.15)$$

which follows directly from Lemma 6.1 (see the Appendix) with $\alpha := p - 1$ and $\beta := 1$. By Remark 5.2, the operator $T := J_\lambda^{\partial g + N_C}$ is firmly nonexpansive and therefore it is a cutter. Moreover, $I - T$ is closed at zero, by the nonexpansivity of T . Hence Conditions 3.1 – 3.4 are satisfied.

6 Appendix

Lemma 6.1 *For any $\alpha, \beta \geq \frac{1}{2}$, there is $c \in (0, 1)$ such that*

$$\left(\sum_{i=1}^n |x_i|^{\alpha+\beta} \right)^2 \geq c \sum_{i=1}^n |x_i|^{2\alpha} \cdot \sum_{i=1}^n |x_i|^{2\beta} \quad (6.1)$$

for all $x \in \mathbb{R}^n$.

Proof. We have to show the following inequality:

$$\left(\|x\|_{\alpha+\beta}^{\alpha+\beta} \right)^2 \geq c \|x\|_{2\alpha}^{2\alpha} \cdot \|x\|_{2\beta}^{2\beta}. \quad (6.2)$$

The norms $\|\cdot\|_{2\alpha}$, $\|\cdot\|_{2\beta}$ and $\|\cdot\|_{\alpha+\beta}$ are all equivalent; hence

$$\|x\|_{\alpha+\beta} \geq c \|x\|_{2\alpha} \quad \text{and} \quad \|x\|_{\alpha+\beta} \geq c \|x\|_{2\beta} \quad (6.3)$$

for some $c > 0$ and all $x \in \mathbb{R}^n$. Without any loss of generality, we can assume that $c \in (0, 1)$. Then

$$\left(\|x\|_{\alpha+\beta}^{\alpha+\beta} \right)^2 = \|x\|_{\alpha+\beta}^{2\alpha} \cdot \|x\|_{\alpha+\beta}^{2\beta} \geq c^{2\alpha+2\beta} \|x\|_{2\alpha}^{2\alpha} \cdot \|x\|_{2\beta}^{2\beta}, \quad (6.4)$$

which yields our assertion. ■

Acknowledgments. The third author was partially supported by Israel Science Foundation (ISF) Grant number 647/07, the Fund for the Promotion of Research at the Technion and by the Technion VPR Fund.

References

- [1] R. Aharoni, A. Berman and Y. Censor, An interior points algorithm for the convex feasibility problem, *Advances in Applied Mathematics* **4** (1983), 479–489.
- [2] A. Auslender, *Optimisation: Méthodes Numériques*, Masson, Paris, 1976.
- [3] H. H. Bauschke, The approximation of fixed points of compositions of nonexpansive mapping in Hilbert space, *Journal of Mathematical Analysis and Applications* **202** (1996), 150–159.
- [4] H. H. Bauschke and J. M. Borwein, On projection algorithms for solving convex feasibility problems, *SIAM Review* **38** (1996), 367–426.
- [5] H. H. Bauschke and P. L. Combettes, A weak-to-strong convergence principle for Fejér-monotone methods in Hilbert spaces, *Mathematics of Operations Research* **26** (2001), 248–264.
- [6] H. H. Bauschke and P. L. Combettes, *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*, Springer, Heidelberg, 2010.
- [7] D. P. Bertsekas and J. N. Tsitsiklis, *Parallel and Distributed Computation: Numerical Methods*, Prentice-Hall International, Englewood Cliffs, NJ, USA, 1989.
- [8] F. E. Browder, Fixed point theorems for noncompact mappings in Hilbert space, *Proceedings of the National Academy of Sciences of the United States of America* **53** (1965), 1272–1276.
- [9] A. Cegielski, A generalization of the Opial theorem, *Control and Cybernetics* **36** (2007), 601–610.
- [10] A. Cegielski, Generalized relaxations of nonexpansive operators and convex feasibility problems, *Contemporary Mathematics* **513** (2010), 111–123.
- [11] A. Cegielski, *Iterative Methods for Fixed Point Problems in Hilbert Spaces*, Lecture Notes in Mathematics **2057**, Springer, Heidelberg, 2012.

- [12] A. Cegielski and Y. Censor, Opial-type theorems and the common fixed point problem, in: H. H. Bauschke, R. S. Burachik, P. L. Combettes, V. Elser, D. R. Luke and H. Wolkowicz (Editors), *Fixed-Point Algorithms for Inverse Problems in Science and Engineering*, Optimization and Its Applications 49, Springer, New York, NY, USA, 2011, 155-183.
- [13] A. Cegielski and R. Zalas, Methods for variational inequality problems over the intersection of fixed point sets of quasi-nonexpansive operators, *Numerical Functional Analysis and Optimization* **34** (2013), 255–283.
- [14] Y. Censor and A. Gibali, Projections onto super-half-spaces for monotone variational inequality problems in finite-dimensional spaces, *Journal of Nonlinear and Convex Analysis* **9** (2008), 461–475.
- [15] Y. Censor and A. Segal, The split common fixed point problem for directed operators, *Journal of Convex Analysis* **16** (2009), 587–600.
- [16] Y. Censor and A. Segal, On the string averaging method for sparse common fixed points problems, *International Transactions in Operational Research* **16** (2009), 481–494.
- [17] P. L. Combettes, Quasi-Fejérian analysis of some optimization algorithms, in: D. Butnariu, Y. Censor and S. Reich (Editors), *Inherently Parallel Algorithms in Feasibility and Optimization and Their Applications*, Elsevier Science Publishers, Amsterdam, The Netherlands, 2001, pp. 115–152.
- [18] G. Crombez, A geometrical look at iterative methods for operators with fixed points, *Numerical Functional Analysis and Optimization* **26** (2005), 157–175.
- [19] G. Crombez, A hierarchical presentation of operators with fixed points on Hilbert spaces, *Numerical Functional Analysis and Optimization* **27** (2006), 259–277.
- [20] F. Facchinei and J. S. Pang, *Finite-Dimensional Variational Inequalities and Complementarity Problems*, Volume I and Volume II, Springer, New York, NY, 2003.
- [21] R. Fletcher, *Practical Methods of Optimization*, John Wiley, Chichester, 1987.

- [22] M. Fukushima, A relaxed projection method for variational inequalities, *Mathematical Programming* **35** (1986), 58–70.
- [23] C. Geiger and Ch. Kanzow, *Numerische Verfahren zur Lösung unrestrictingierter Optimierungsaufgaben*, Springer, Berlin, 1999.
- [24] A. Gibali, Investigation of Iterative Algorithms for Solving Variational Inequality Problems, M.Sc. Thesis, University of Haifa, November 2007.
- [25] J.-B. Hiriart-Urruty and C. Lemaréchal, *Convex Analysis and Minimization Algorithms*, Vol I, Vol II, Springer, Berlin, 1993.
- [26] S. A. Hirstoaga, Iterative selection methods for common fixed point problems, *Journal of Mathematical Analysis and Applications* **324** (2006), 1020–1035.
- [27] D. Kinderlehrer and G. Stampacchia, *An Introduction to Variational Inequalities and Their Applications*, Academic Press, New York, 1980.
- [28] Șt. Mărușter, Quasi-nonexpansivity and the convex feasibility problem, *Analele Științifice ale Universității “Alexandru Ioan Cuza” din Iași Informatică* **15** (2005), 47–56.
- [29] Ș. Mărușter and C. Popirlan, On the Mann-type iteration and the convex feasibility problem, *Journal of Computational and Applied Mathematics* **212** (2008), 390–396.
- [30] M. A. Noor, Some developments in general variational inequalities, *Applied Mathematics and Computation* **152** (2004), 197–277.
- [31] H. F. Senter and W. G. Dotson, Jr., Approximating fixed points of non-expansive mappings, *Proceedings of the American Mathematical Society* **44** (1974), 375–380.
- [32] N. Xiu and J. Zhang, Some recent advances in projection-type methods for variational inequalities, *Journal of Computational and Applied Mathematics* **152** (2003), 559–585.
- [33] I. Yamada and N. Ogura, Adaptive projected subgradient method for asymptotic minimization of sequence of nonnegative convex operators, *Numerical Functional Analysis and Optimization* **25** (2004), 593–617.

- [34] I. Yamada and N. Ogura, Hybrid steepest descent method for variational inequality problem over the fixed point set of certain quasi-nonexpansive mappings, *Numerical Functional Analysis And Optimization* **25** (2004), 619-655.
- [35] I. Yamada, M. Yukawa and M. Yamagishi, Minimizing the Moreau envelope of nonsmooth convex functions over the fixed point set of certain quasinonexpansive mapping, In: H. H. Basuchke, R. S. Burachik, P. L. Combettes, V. Elser, D. R. Luke, H. Wolkowicz (Editors), *Fixed-point algorithms for inverse problems in science and engineering*, Optimization and Its Applications 49, Springer, New York, NY, USA, 2011, 345–390.
- [36] M. Zaknoon, Algorithmic developments for the convex feasibility problem, Ph.D. Thesis, University of Haifa, 2003.